



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT


KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-646
Date- 25/08/2026

Cuttack District Block Wise Agro-met Advisory Service Bulletin

Tuesday, 25th August, 2026
(For the period 25th to 29th August, 2026)



Issued by
District Agromet Unit,
KrishiVigyan Kendra Cuttack



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightning alert in your area






GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-646
Date- 25/08/2026

STANDARD PRECIPITATION INDEX AND NDVI MAP



LEGEND (Medium range block level rainfall forecast)		
	No Rainfall	00 mm
	Very light rainfall	01-05 mm
	Light rainfall	06-10 mm
	Light to moderate rainfall	11-30 mm
	Moderate to heavy rainfall	31-65 mm
	Heavy to very heavy rainfall	66-115 mm
	Extremely heavy rainfall	>116 mm

DrRanjan Kumar Mohanta
 (Nodal officer)

**Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area**





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-646
Date- 25/08/2026

Weather (Past and Forecasted) based Disease and pest in Major Crops and vegetables

Sl No	Crop	Stage and condition of the crop	Anticipated Disease and pest	
1	Rice	Tillering		Blast
				Leaf streak
				Stem Borer
				Thrips
				Gall midge
2	Sugarcane	Vegetative to Tillering		Red rot
				Wilt
				Early shoot borer
3	Maize	5 week stage		Stem fly
				Leaf blight
				Stalk rot



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-646
Date- 25/08/2026

3	Jute	30-50days		Stem rot
				Indigo catterpillar
4	Brinjal, tomoto, chilli	seedling		Damping off
				Stem rot
5	Tomato and brinjal			Mealy bug
				Mites
				Bacterial wilt
				White fly
				Jassid
6	okra	Fruting		cercospora leaf spot
7	Banana			Sigatoka
8	Papaya			Yellow mosaic virus



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-646
Date- 25/08/2026

AGROMET ADVISORY BULLETIN FOR ATHAGAD

District Past week average weather condition and Block past week average Rainfall condition

The block received average 16 mm and district received average 08 mm rainfall during the last week. The mean daily maximum and minimum temperature was 32.34°C (+0.34) and 23.62°C (-0.5), respectively. The mean relative humidity was 83-88%.

Forecasted period: Today(8.30IST) to 30th August (8.30IST)2026

Weather advisory: Light to Moderate rainfall/thundershower is forecasted for today to Saturday in the block. The day temperature will be 32-33°C and night temperature will be 25°C for upcoming 5 days. Maximum and minimum relative humidity will be 38-90% and 49-68% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Athagad (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	25/08/2026	26/08/2026	27/08/2026	28/08/2026	29/08/2026
Rainfall (mm)	7	7	6	15	25
T-Max (°C)	32	33	33	33	33
T-Min (°C)	25	25	25	25	25
Cloud Cover (octa)	7	5	6	8	8
RH Max (%)	38	38	88	80	90
RH Min (%)	60	60	56	49	68
Wind Speed (kmph)	9	8	6	8	10
Wind Direction (deg)	168	243	215	176	138

Agromet Advisory

General advisory: Farmers should drain out excess water from vegetable and rice field. They may go for sowing, fertilizer, pesticide application, intercultural operations & human labour engagement before Friday based upon rainfall forecast.

Kharif Rice:

- The crop is at tillering stage. Maintain about 2-3 cm standing water during tillering stage. Problem of weeds may aggravate after transplanting, apply Bensulfuron methyl 0.6% + Pretilachlor 6% GR with sand mix at 1:1 ratio @ 4 kg/acre within 3-7 days after transplanting for controlling weeds.
- If **leaf blast** is observed in nursery, spray Tebuconazole 50% + Trifloxystrobin 25% WG @ 0.4 g or Isoprothiolane 40EC @ 1.5 ml per litre of water. Repeat the spray at 7-10 days interval.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre
- Weekly Temperature below 28deg c and continuous rainfall lead to infestation of **stem borer** at tillering stage of rice, to control this apply Azadirachtin 0.15% EC @ 800 ml/acre or Cartap hydrochloride 4G @ 10kg/acre.
- Due to forecasted slight increase in temperature there may be increase in infestation of **thrips**, to control this Spray Lambda-cyhalothrin 5% EC @ 2 ml/ litre water or Thiamethoxam 25% WG @ 5g/ 15 litre water.
- Due to slight increase temperature and high humidity there will be chances of infestation of **bacterial leaf streak**, to control apply Plantomycin @ 1 g/litre along with Copper oxychloride @ 1g/litre of water using 200 litre of water per acre.

Maize: Maize crop is highly sensitive to waterlogged conditions during seedling stage. Hence, drain out excess water from the field as early as possible.

- After cessation of rains apply booster dose of fertilizer.
- Weeds may be controlled after rainfall.
- Stem fly** in Maize infects the plants from sowing to 5 weeks after sowing, temperature between 25-30 deg. celsius and humidity above 60% favour pest incidence. In vegetative stage to control stem fly spray Methyl demeton @ 1 ml/ltr or Dimethoate @ 2ml/ltr or apply Carbofuran 3G granules @ 14 kg/acre.
- High humid conditions, temperature between 18-27 Deg. Celsius favours the occurrence of **leaf blight** in Maize. To control spray Azoxystrobin + Mancozeb 300 g /ac. Due to water logging condition stem rot may be found, then farmers should apply carbendazim @ 2g/ltr water.

VEGETABLES:

Anticipation of decrease in rainfall or less cloudy weather lead to infestation of **Mealy bug and **mites** in vegetables. Take up spraying of profenophos 50% EC @ 400ml/ac against mealy bug and propargite 57%EC/ fenazaquin 10%EC / Fenpyroximate 5% EC @ 400ml/ac against mites which are more prevalent in dry weather.**

- Due to past rain, Chances of **fruit rot** infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary, spray second and third round at 15 days interval.
- If incidence of **Damping off** is noticed in Tomato, Brinjal, Chilli nurseries, drench nursery bed with Copper-Oxy-Chloride @ 3 g per liter of water.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever **sucking pests** incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. To prevent of soil borne diseases like **seedling rot, stem rot and wilt** diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control **Cercospora leaf spot**.

Fruits: Spray mancozeb 2g/litre, to control sigatoka leaf spot in **banana**. To control **papaya yellow mosaic virus**, spray Imidachloprid @ 0.5 ml / litre.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-646
Date- 25/08/2026

AGROMET ADVISORY BULLETIN FOR BADAMBA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 03 mm and district received average 08 mm rainfall during the last week. The mean daily maximum and minimum temperature was 32.34°C (+0.34) and 23.62°C (-0.5), respectively. The mean relative humidity was 83-88%.

Forecasted period: Today(8.30IST) to 30th August (8.30IST)2026

Weather advisory *Light to Moderate rainfall/thundershower is forecasted for today to Saturday in the block.* The day temperature will be 32-33°C and night temperature will be 25°C for upcoming 5 days. Maximum and minimum relative humidity will be 37-90% and 49-68% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Badamba (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	25/08/2026	26/08/2026	27/08/2026	28/08/2026	29/08/2026
Rainfall(mm)	8	7	6	15	26
T-Max(°C)	32	32	33	33	32
T-Min(°C)	25	25	25	25	25
Cloud Cover(octa)	7	6	6	7	8
RH Max (%)	37	37	80	80	90
RH Min (%)	60	60	56	49	68
Wind Speed (Kmph)	9	8	6	8	10
Wind Direction (deg)	168	243	215	176	138

Agromet Advisory

General advisory: Farmers should drain out excess water from vegetable and rice field. They may go for sowing, fertilizer, pesticide application, intercultural operations & human labour engagement before Friday based upon rainfall forecast.

Kharif Rice:

- The crop is at tillering stage. Maintain about 2-3 cm standing water during tillering stage. Problem of weeds may aggravate after transplanting, apply Bensulfuron methyl 0.6% + Pretilachlor 6% GR with sand mix at 1:1 ratio @ 4 kg/acre within 3-7 days after transplanting for controlling weeds.
- If **leaf blast** is observed in nursery, spray Tebuconazole 50% + Trifloxystrobin 25% WG @ 0.4 g or Isoprothiolane 40EC @ 1.5 ml per litre of water. Repeat the spray at 7-10 days interval.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre
- Weekly Temperature below 28deg c and continuous rainfall lead to infestation of **stem borer** at tillering stage of rice, to control this apply Azadirachtin 0.15% EC @ 800 ml/acre or Cartap hydrochloride 4G @ 10kg/acre.
- Due to forecasted slight increase in temperature there may be increase in infestation of **thrips**, to control this Spray Lambda-cyhalothrin 5% EC @ 2 ml/ litre water or Thiamethoxam 25% WG @ 5g/ 15 litre water.
- Due to slight increase temperature and high humidity there will be chances of infestation of **bacterial leaf streak**, to control apply Plantomycin @ 1 g/litre along with Copper oxychloride @ 1g/litre of water using 200 litre of water per acre.

Maize: Maize crop is highly sensitive to waterlogged conditions during seedling stage. Hence, drain out excess water from the field as early as possible.

- After cessation of rains apply booster dose of fertilizer.
- Weeds may be controlled after rainfall.
- Stem fly** in Maize infects the plants from sowing to 5 weeks after sowing, temperature between 25-30 deg. celsius and humidity above 60% favour pest incidence. In vegetative stage to control stem fly spray Methyl demeton @ 1 ml/ltr or Dimethoate @ 2ml/ltr or apply Carbofuran 3G granuals @ 14 kg/acre.
- High humid conditions, temperature between 18-27 Deg. Celsius favours the occurrence of **leaf blight** in Maize. To control spray Azoxystrobin + Mancozeb 300 g /ac. Due to water logging condition stem rot may be found, then farmers should apply carbendazim @ 2g/ltr water.

VEGETABLES:

Anticipation of decrease in rainfall or less cloudy weather lead to infestation of **Mealy bug** and **mites** in vegetables. Take up spraying of profenophos 50% EC @ 400ml/ac against mealy bug and propargite 57%EC/ fenazaquin 10%EC / Fenpyroximate 5% EC @ 400ml/ac against mites which are more prevalent in dry weather.

- Due to past rain, Chances of **fruit rot** infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary, spray second and third round at 15 days interval.
- If incidence of **Damping off** is noticed in Tomato, Brinjal, Chilli nurseries, drench nursery bed with Copper-Oxy-Chloride @ 3 g per liter of water.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever **sucking pests** incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. To prevent of soil borne diseases like **seedling rot, stem rot and wilt** diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control **Cercospora leaf spot**.

Fruits: Spray mancozeb 2g/litre, to control sigatoka leaf spot in banana. To control papaya yellow mosaic virus, spray Imidachloprid @ 0.5 ml / litre.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-646
Date- 25/08/2026

AGROMET ADVISORY BULLETIN FOR BANKI

District Past week average weather condition

The district received average 08 mm rainfall during the last week. The mean daily maximum and minimum temperature was 32.34°C (+0.34) and 23.62°C (-0.5), respectively. The mean relative humidity was 83-88%.

Forecasted period: Today(8.30IST) to 30th August (8.30IST)2026

Weather advisory: *Light to Moderate rainfall/thundershower is forecasted for today to Saturday in the block.* The day temperature will be 32-33°C and night temperature will be 25°C for upcoming 5 days. Maximum and minimum relative humidity will be 37-90% and 49-68% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Banki (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	25/08/2026	26/08/2026	27/08/2026	28/08/2026	29/08/2026
Rainfall(mm)	8	7	6	15	26
T-Max(°C)	32	32	33	33	32
T-Min(°C)	25	25	25	25	25
Cloud Cover(octa)	7	6	6	7	8
RH Max (%)	37	37	80	80	90
RH Min (%)	60	60	56	49	68
Wind Speed (Kmph)	9	8	6	8	10
Wind Direction (deg)	168	243	215	176	138

Agromet Advisory

General advisory: Farmers should drain out excess water from vegetable and rice field. They may go for sowing, fertilizer, pesticide application, intercultural operations & human labour engagement before Friday based upon rainfall forecast.

Kharif Rice:

- The crop is at tillering stage. Maintain about 2-3 cm standing water during tillering stage. Problem of weeds may aggravate after transplanting, apply Bensulfuron methyl 0.6% + Pretilachlor 6% GR with sand mix at 1:1 ratio @ 4 kg/acre within 3-7 days after transplanting for controlling weeds.
- If **leaf blast** is observed in nursery, spray Tebuconazole 50% + Trifloxystrobin 25% WG @ 0.4 g or Isoprothiolane 40EC @ 1.5 ml per litre of water. Repeat the spray at 7-10 days interval.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre
- Weekly Temperature below 28deg c and continuous rainfall lead to infestation of **stem borer** at tillering stage of rice, to control this apply Azadirachtin 0.15% EC @ 800 ml/acre or Cartap hydrochloride 4G @ 10kg/acre.
- Due to forecasted slight increase in temperature there may be increase in infestation of **thrips**, to control this Spray Lambda-cyhalothrin 5% EC @ 2 ml/ litre water or Thiamethoxam 25% WG @ 5g/ 15 litre water.
- Due to slight increase temperature and high humidity there will be chances of infestation of **bacterial leaf streak**, to control apply Plantomycin @ 1 g/litre along with Copper oxychloride @ 1g/litre of water using 200 litre of water per acre.

Maize: Maize crop is highly sensitive to waterlogged conditions during seedling stage. Hence, drain out excess water from the field as early as possible.

- After cessation of rains apply booster dose of fertilizer.
- Weeds may be controlled after rainfall.
- Stem fly** in Maize infects the plants from sowing to 5 weeks after sowing, temperature between 25-30 deg. celsius and humidity above 60% favour pest incidence. In vegetative stage to control stem fly spray Methyl demeton @ 1 ml/ltr or Dimethoate @ 2ml/ltr or apply Carbofuran 3G granules @ 14 kg/acre.
- High humid conditions, temperature between 18-27 Deg. Celsius favours the occurrence of **leaf blight** in Maize. To control spray Azoxystrobin + Mancozeb 300 g /ac. Due to water logging condition stem rot may be found, then farmers should apply carbendazim @ 2g/ltr water.

VEGETABLES:

Anticipation of decrease in rainfall or less cloudy weather lead to infestation of **Mealy bug** and **mites** in vegetables. Take up spraying of profenophos 50% EC @ 400ml/ac against mealy bug and propargite 57%EC/ fenazaquin 10%EC / Fenpyroximate 5% EC @ 400ml/ac against mites which are more prevalent in dry weather.

- Due to past rain, Chances of **fruit rot** infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary, spray second and third round at 15 days interval.
- If incidence of **Damping off** is noticed in Tomato, Brinjal, Chilli nurseries, drench nursery bed with Copper-Oxy-Chloride @ 3 g per liter of water.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever **sucking pests** incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. To prevent of soil borne diseases like **seedling rot, stem rot and wilt** diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control **Cercospora leaf spot**.

Fruits: Spray mancozeb 2g/litre, to control sigatoka leaf spot in banana. To control papaya yellow mosaic virus, spray Imidachloprid @ 0.5 ml / litre.



Dr Ranjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-646
Date- 25/08/2026

AGROMET ADVISORY BULLETIN FOR BANKI-DAMPADA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 09 mm and district received average 08 mm rainfall during the last week. The mean daily maximum and minimum temperature was 32.34°C (+0.34) and 23.62°C (-0.5), respectively. The mean relative humidity was 83-88%.

Forecasted period: Today(8.30IST) to 30th August (8.30IST)2026

Weather Advisory:: *Light to Moderate rainfall/thundershower is forecasted for today to Saturday in the block.* The day temperature will be 32-33°C and night temperature will be 25°C for upcoming 5 days. Maximum and minimum relative humidity will be 88-91% and 50-68% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Banki-Dampada (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	25/08/2026	26/08/2026	27/08/2026	28/08/2026	29/08/2026
Rainfall(mm)	7	7	6	15	26
T-Max(°C)	32	33	33	33	33
T-Min(°C)	25	25	25	25	25
Cloud Cover(octa)	7	5	6	7	8
RH Max (%)	91	89	88	89	91
RH Min (%)	60	63	58	50	68
Wind Speed (Kmph)	9	8	7	7	9
Wind Direction (deg)	198	255	244	189	145

Agromet Advisory

General advisory: Farmers should drain out excess water from vegetable and rice field. They may go for sowing, fertilizer, pesticide application, intercultural operations & human labour engagement before Friday based upon rainfall forecast.

Kharif Rice:

- The crop is at tillering stage. Maintain about 2-3 cm standing water during tillering stage. Problem of weeds may aggravate after transplanting, apply Bensulfuron methyl 0.6% + Pretilachlor 6% GR with sand mix at 1:1 ratio @ 4 kg/acre within 3-7 days after transplanting for controlling weeds.
- If **leaf blast** is observed in nursery, spray Tebuconazole 50% + Trifloxystrobin 25% WG @ 0.4 g or Isoprothiolane 40EC @ 1.5 ml per litre of water. Repeat the spray at 7-10 days interval.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre
- Weekly Temperature below 28deg c and continuous rainfall lead to infestation of **stem borer** at tillering stage of rice, to control this apply Azadirachtin 0.15% EC @ 800 ml/acre or Cartap hydrochloride 4G @ 10kg/acre.
- Due to forecasted slight increase in temperature there may be increase in infestation of **thrips**, to control this Spray Lambda-cyhalothrin 5% EC @ 2 ml/ litre water or Thiamethoxam 25% WG @ 5g/ 15 litre water.
- Due to slight increase temperature and high humidity there will be chances of infestation of **bacterial leaf streak**, to control apply Plantomycin @ 1 g/litre along with Copper oxychloride @ 1g/litre of water using 200 litre of water per acre.

Maize: Maize crop is highly sensitive to waterlogged conditions during seedling stage. Hence, drain out excess water from the field as early as possible.

- After cessation of rains apply booster dose of fertilizer.
- Weeds may be controlled after rainfall.
- Stem fly** in Maize infects the plants from sowing to 5 weeks after sowing, temperature between 25-30 deg. celsius and humidity above 60% favour pest incidence. In vegetative stage to control stem fly spray Methyl demeton @ 1 ml/ltr or Dimethoate @ 2ml/ltr or apply Carbofuran 3G granuals @ 14 kg/acre.
- High humid conditions, temperature between 18-27 Deg. Celsius favours the occurrence of **leaf blight** in Maize. To control spray Azoxystrobin + Mancozeb 300 g /ac. Due to water logging condition stem rot may be found, then farmers should apply carbendazim @ 2g/ltr water.

VEGETABLES:

Anticipation of decrease in rainfall or less cloudy weather lead to infestation of **Mealy bug** and **mites** in vegetables. Take up spraying of profenophos 50% EC @ 400ml/ac against mealy bug and propargite 57%EC/ fenazaquin 10%EC / Fenpyroximate 5% EC @ 400ml/ac against mites which are more prevalent in dry weather.

- Due to past rain, Chances of **fruit rot** infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary, spray second and third round at 15 days interval.
- If incidence of **Damping off** is noticed in Tomato, Brinjal, Chilli nurseries, drench nursery bed with Copper-Oxy-Chloride @ 3 g per liter of water.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever **sucking pests** incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. To prevent of soil borne diseases like **seedling rot, stem rot and wilt** diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control **Cercospora leaf spot**.

Fruits: Spray mancozeb 2g/litre, to control sigatoka leaf spot in banana. To control papaya yellow mosaic virus, spray Imidachloprid @ 0.5 ml / litre.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-646
Date- 25/08/2026

AGROMET ADVISORY BULLETIN FOR BARANGA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 07 mm and district received average 08 mm rainfall during the last week. The mean daily maximum and minimum temperature was 32.34°C (+0.34) and 23.62°C (-0.5), respectively. The mean relative humidity was 83-88%.

Forecasted period: Today(8.30IST) to 30th August (8.30IST)2026

Weather advisory: *Light to Moderate rainfall/thundershower is forecasted for today to Saturday in the block.* The day temperature will be 33-34°C and night temperature will be 25°C for upcoming 5 days. Maximum and minimum relative humidity will be 88-91% and 50-68% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Baranga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	25/08/2026	26/08/2026	27/08/2026	28/08/2026	29/08/2026
Rainfall(mm)	8	7	6	15	25
T-Max(°C)	33	33	34	34	33
T-Min(°C)	25	25	25	25	25
Cloud Cover	7	5	6	7	8
RH Max (%)	91	89	88	89	91
RH Min (%)	60	63	58	50	68
Wind Speed (Kmph)	9	8	7	7	9
Wind Direction (deg)	198	255	244	189	145

Agromet Advisory

General advisory: Farmers should drain out excess water from vegetable and rice field. They may go for sowing, fertilizer, pesticide application, intercultural operations & human labour engagement before Friday based upon rainfall forecast.

Kharif Rice:

- The crop is at tillering stage. Maintain about 2-3 cm standing water during tillering stage. Problem of weeds may aggravate after transplanting, apply Bensulfuron methyl 0.6% + Pretilachlor 6% GR with sand mix at 1:1 ratio @ 4 kg/acre within 3-7 days after transplanting for controlling weeds.
- If leaf blast is observed in nursery, spray Tebuconazole 50% + Trifloxystrobin 25% WG @ 0.4 g or Isoprothiolane 40EC @ 1.5 ml per litre of water. Repeat the spray at 7-10 days interval.
- In Gall midge affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre
- Weekly Temperature below 28deg c and continuous rainfall lead to infestation of stem borer at tillering stage of rice, to control this apply Azadirachtin 0.15% EC @ 800 ml/acre or Cartap hydrochloride 4G @ 10kg/acre.
- Due to forecasted slight increase in temperature there may be increase in infestation of thrips, to control this Spray Lambda-cyhalothrin 5% EC @ 2 ml/ litre water or Thiamethoxam 25% WG @ 5g/ 15 litre water.
- Due to slight increase temperature and high humidity there will be chances of infestation of bacterial leaf streak, to control apply Plantomycin @ 1 g/litre along with Copper oxychloride @ 1g/litre of water using 200 litre of water per acre.

Maize: Maize crop is highly sensitive to waterlogged conditions during seedling stage. Hence, drain out excess water from the field as early as possible.

- After cessation of rains apply booster dose of fertilizer.
- Weeds may be controlled after rainfall.
- Stem fly in Maize infects the plants from sowing to 5 weeks after sowing, temperature between 25-30 deg. celsius and humidity above 60% favour pest incidence. In vegetative stage to control stem fly spray Methyl demeton @ 1 ml/lit or Dimethoate @ 2ml/lit or apply Carbofuran 3G granules @ 14 kg/acre.
- High humid conditions, temperature between 18-27 Deg. Celsius favours the occurrence of leaf blight in Maize. To control spray Azoxystrobin + Mancozeb 300 g /ac. Due to water logging condition stem rot may be found, then farmers should apply carbendazim @ 2g/ltr water.

VEGETABLES:

Anticipation of decrease in rainfall or less cloudy weather lead to infestation of Mealy bug and mites in vegetables. Take up spraying of profenophos 50% EC @ 400ml/ac against mealy bug and propargite 57%EC/ fenazaquin 10%EC / Fenpyroximate 5% EC @ 400ml/ac against mites which are more prevalent in dry weather.

- Due to past rain, Chances of fruit rot infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary, spray second and third round at 15 days interval.
- If incidence of Damping off is noticed in Tomato, Brinjal, Chilli nurseries, drench nursery bed with Copper-Oxy-Chloride @ 3 g per liter of water.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever sucking pests incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. To prevent of soil borne diseases like seedling rot, stem rot and wilt diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control Cercospora leaf spot.

Fruits: Spray mancozeb 2g/litre, to control sigatoka leaf spot in banana. To control papaya yellow mosaic virus, spray Imidachloprid @ 0.5 ml / litre.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-646
Date- 25/08/2026

AGROMET ADVISORY BULLETIN FOR CUTTACK SADAR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 10 mm and district received average 08 mm rainfall during the last week. The mean daily maximum and minimum temperature was 32.34°C (+0.34) and 23.62°C (-0.5), respectively. The mean relative humidity was 83-88%.

Forecasted period: Today(8.30IST) to 30th August (8.30IST)2026

Weather advisory: *Light to Moderate rainfall/thundershower is forecasted for today to Saturday in the block.* The day temperature will be 33-34°C and night temperature will be 25°C for upcoming 5 days. Maximum and minimum relative humidity will be 88-91% and 50-68% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Cuttack sadar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	25/08/2026	26/08/2026	27/08/2026	28/08/2026	29/08/2026
Rainfall(mm)	8	7	6	16	24
T-Max(°C)	33	34	34	34	33
T-Min(°C)	25	25	25	25	25
Cloud Cover(octa)	7	6	7	8	8
RH Max (%)	91	89	88	89	91
RH Min (%)	60	63	58	50	68
Wind Speed (Kmph)	9	8	7	7	9
Wind Direction (deg)	198	255	244	189	145

Agromet Advisory

General advisory: Farmers should drain out excess water from vegetable and rice field. They may go for sowing, fertilizer, pesticide application, intercultural operations & human labour engagement before Friday based upon rainfall forecast.

Kharif Rice:

- The crop is at tillering stage. Maintain about 2-3 cm standing water during tillering stage. Problem of weeds may aggravate after transplanting, apply Bensulfuron methyl 0.6% + Pretilachlor 6% GR with sand mix at 1:1 ratio @ 4 kg/acre within 3-7 days after transplanting for controlling weeds.
- If leaf blast is observed in nursery, spray Tebuconazole 50% + Trifloxystrobin 25% WG @ 0.4 g or Isoprothiolane 40EC @ 1.5 ml per litre of water. Repeat the spray at 7-10 days interval.
- In Gall midge affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre
- Weekly Temperature below 28deg c and continuous rainfall lead to infestation of stem borer at tillering stage of rice, to control this apply Azadirachtin 0.15% EC @ 800 ml/acre or Cartap hydrochloride 4G @ 10kg/acre.
- Due to forecasted slight increase in temperature there may be increase in infestation of thrips, to control this Spray Lambda-cyhalothrin 5% EC @ 2 ml/ litre water or Thiamethoxam 25% WG @ 5g/ 15 litre water.
- Due to slight increase temperature and high humidity there will be chances of infestation of bacterial leaf streak, to control apply Plantomycin @ 1 g/litre along with Copper oxychloride @ 1g/litre of water using 200 litre of water per acre.

Sugarcane:

- Drain out excess water from the field as early as possible.
- After cessation of rains weeding and earthing up should be done in this time after that Top dressing of Urea @30 kg/acre for 30-60 days old sugarcane plant.
- For Red rot and wilt disease, spray the crop with Mancozeb 0.3% or carbendazim 0.1% or Azoxystrobin 18.2%SC/ Difenconazole 11.45SC @ 1ml per liter water in dry days.
- For early shoot borer Spray Chloropyriphos @ 400 ml/ac in 200 lts of water after cessation of rain.

VEGETABLES:

Anticipation of decrease in rainfall or less cloudy weather lead to infestation of Mealy bug and mites in vegetables. Take up spraying of profenophos 50% EC @ 400ml/ac against mealy bug and propargite 57%EC/ fenazaquin 10%EC / Fenpyroximate 5% EC @ 400ml/ac against mites which are more prevalent in dry weather.

- Due to past rain, Chances of fruit rot infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary, spray second and third round at 15 days interval.
- If incidence of Damping off is noticed in Tomato, Brinjal, Chilli nurseries, drench nursery bed with Copper-Oxy-Chloride @ 3 g per liter of water.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever sucking pests incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. To prevent of soil borne diseases like seedling rot, stem rot and wilt diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control Cercospora leaf spot.

Fruits: Spray mancozeb 2g/litre, to control sigatoka leaf spot in banana. To control papaya yellow mosaic virus, spray Imidachloprid @ 0.5 ml / litre.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-646
Date- 25/08/2026

AGROMET ADVISORY BULLETIN FOR KANTAPADA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 08 mm and district received average 08 mm rainfall during the last week. The mean daily maximum and minimum temperature was 32.34°C (+0.34) and 23.62°C (-0.5), respectively. The mean relative humidity was 83-88%.

Forecasted period: Today(8.30IST) to 30th August (8.30IST)2026

Weather advisory: *Light to Moderate rainfall/thundershower is forecasted for today to Saturday in the block.* The day temperature will be 33-34°C and night temperature will be 25-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 85-91% and 50-68% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Kantapada (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	25/08/2026	26/08/2026	27/08/2026	28/08/2026	29/08/2026
Rainfall(mm)	8	7	6	15	24
T-Max(°C)	33	33	34	34	33
T-Min(°C)	25	25	26	26	26
Cloud Cover(octa)	7	5	7	7	8
RH Max (%)	85	89	88	89	91
RH Min (%)	60	63	58	50	68
Wind Speed (Kmph)	9	8	7	7	9
Wind Direction (deg)	198	255	244	189	145

Agromet Advisory

General advisory: Farmers should drain out excess water from vegetable and rice field. They may go for sowing, fertilizer, pesticide application, intercultural operations & human labour engagement before Friday based upon rainfall forecast.

Kharif Rice:

- The crop is at tillering stage. Maintain about 2-3 cm standing water during tillering stage. Problem of weeds may aggravate after transplanting, apply Bensulfuron methyl 0.6% + Pretilachlor 6% GR with sand mix at 1:1 ratio @ 4 kg/acre within 3-7 days after transplanting for controlling weeds.
- If **leaf blast** is observed in nursery, spray Tebuconazole 50% + Trifloxystrobin 25% WG @ 0.4 g or Isoprothiolane 40EC @ 1.5 ml per litre of water. Repeat the spray at 7-10 days interval.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre
- Weekly Temperature below 28deg c and continuous rainfall lead to infestation of **stem borer** at tillering stage of rice, to control this apply Azadirachtin 0.15% EC @ 800 ml/acre or Cartap hydrochloride 4G @ 10kg/acre.
- Due to forecasted slight increase in temperature there may be increase in infestation of **thrips**, to control this Spray Lambda-cyhalothrin 5% EC @ 2 ml/ litre water or Thiamethoxam 25% WG @ 5g/ 15 litre water.
- Due to slight increase temperature and high humidity there will be chances of infestation of **bacterial leaf streak**, to control apply Plantomycin @ 1 g/litre along with Copper oxychloride @ 1g/litre of water using 200 litre of water per acre.

Sugarcane:

- Drain out excess water from the field as early as possible.
- After cessation of rains weeding and earthing up should be done in this time after that Top dressing of Urea @30 kg/acre for 30-60 days old sugarcane plant.
- For **Red rot and wilt disease**, spray the crop with Mancozeb 0.3% or carbendazim 0.1% or Azoxystrobin 18.2%SC/ Difenconazole 11.45SC @ 1ml per liter water in dry days.
- For early shoot borer Spray Chloropyrifos @ 400 ml/ac in 200 lts of water after cessation of rain.

VEGETABLES:

Anticipation of decrease in rainfall or less cloudy weather lead to infestation of **Mealy bug** and **mites** in vegetables. Take up spraying of profenophos 50% EC @ 400ml/ac against mealy bug and propargite 57%EC/ fenazaquin 10%EC / Fenpyroximate 5% EC @ 400ml/ac against mites which are more prevalent in dry weather.

- Due to past rain, Chances of **fruit rot** infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary, spray second and third round at 15 days interval.
- If incidence of **Damping off** is noticed in Tomato, Brinjal, Chilli nurseries, drench nursery bed with Copper-Oxy-Chloride @ 3 g per liter of water.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever **sucking pests** incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. To prevent of soil borne diseases like **seedling rot, stem rot and wilt** diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control **Cercospora leaf spot**.

Fruits:

Spray mancozeb 2g/litre, to control sigatoka leaf spot in **banana**. To control **papaya yellow mosaic virus**, spray Imidachloprid @ 0.5 ml / litre.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-646
Date- 25/08/2026

AGROMET ADVISORY BULLETIN FOR MAHANGA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 03 mm and district received average 08 mm rainfall during the last week. The mean daily maximum and minimum temperature was 32.34°C (+0.34) and 23.62°C (-0.5), respectively. The mean relative humidity was 83-88%.

Forecasted period: Today(8.30IST) to 30th August (8.30IST)2026

Weather advisory: *Light to Moderate rainfall/thundershower is forecasted for today to Saturday in the block.* The day temperature will be and it will be 32-33°C and night temperature will be 25-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 85-91% and 50-68 respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Mahanga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	25/08/2026	26/08/2026	27/08/2026	28/08/2026	29/08/2026
Rainfall(mm)	7	6	6	14	24
T-Max(°C)	32	33	33	33	33
T-Min(°C)	25	25	26	26	26
Cloud Cover(octal)	7	6	6	7	8
RH Max (%)	85	89	88	89	91
RH Min (%)	60	63	58	50	68
Wind Speed (Kmph)	9	8	7	7	9
Wind Direction (deg)	198	255	244	189	145

Agromet Advisory

General advisory: *Farmers should drain out excess water from vegetable and rice field. They may go for sowing, fertilizer, pesticide application, intercultural operations & human labour engagement before Friday based upon rainfall forecast.*

Kharif Rice:

- The crop is at tillering stage. Maintain about 2-3 cm standing water during tillering stage. Problem of weeds may aggravate after transplanting, apply Bensulfuron methyl 0.6% + Pretilachlor 6% GR with sand mix at 1:1 ratio @ 4 kg/acre within 3-7 days after transplanting for controlling weeds.
- If **leaf blast** is observed in nursery, spray Tebuconazole 50% + Trifloxystrobin 25% WG @ 0.4 g or Isoprothiolane 40EC @ 1.5 ml per litre of water. Repeat the spray at 7-10 days interval.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre
- Weekly Temperature below 28deg c and continuous rainfall lead to infestation of **stem borer** at tillering stage of rice, to control this apply Azadirachtin 0.15% EC @ 800 ml/acre or Cartap hydrochloride 4G @ 10kg/acre.
- Due to forecasted slight increase in temperature there may be increase in infestation of **thrips**, to control this Spray Lambda-cyhalothrin 5% EC @ 2 ml/ litre water or Thiamethoxam 25% WG @ 5g/ 15 litre water.
- Due to slight increase temperature and high humidity there will be chances of infestation of **bacterial leaf streak**, to control apply Plantomycin @ 1 g/litre along with Copper oxychloride @ 1g/litre of water using 200 litre of water per acre.

Sugarcane:

- Drain out excess water from the field as early as possible.
- After cessation of rains weeding and earthing up should be done in this time after that Top dressing of Urea @ 30 kg/acre for 30-60 days old sugarcane plant.
- For **Red rot and wilt disease**, spray the crop with Mancozeb 0.3% or carbendazim 0.1% or Azoxystrobin 18.2% SC/ Difenconazole 11.45SC @ 1ml per liter water in dry days.
- For early shoot borer Spray Chloropyrifos @ 400 ml/ac in 200 lts of water after cessation of rain.

VEGETABLES:

Anticipation of decrease in rainfall or less cloudy weather lead to infestation of **Mealy bug** and **mites** in vegetables. Take up spraying of profenophos 50% EC @ 400ml/ac against mealy bug and propargite 57% EC/ fenazaquin 10% EC / Fenpyroximate 5% EC @ 400ml/ac against mites which are more prevalent in dry weather.

- Due to past rain, Chances of **fruit rot** infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary, spray second and third round at 15 days interval.
- If incidence of **Damping off** is noticed in Tomato, Brinjal, Chilli nurseries, drench nursery bed with Copper-Oxy-Chloride @ 3 g per liter of water.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever **sucking pests** incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. To prevent of soil borne diseases like **seedling rot, stem rot and wilt** diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control **Cercospora leaf spot**.

Fruits: Spray mancozeb 2g/litre, to control sigatoka leaf spot in banana. To control papaya yellow mosaic virus, spray Imidachloprid @ 0.5 ml / litre.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-646
Date- 25/08/2026

AGROMET ADVISORY BULLETIN FOR NARASINGHPUR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 06 mm and district received average 08 mm rainfall during the last week. The mean daily maximum and minimum temperature was 32.34°C (+0.34) and 23.62°C (-0.5), respectively. The mean relative humidity was 83-88%.

Forecasted period: Today(8.30IST) to 30th August (8.30IST)2026

Weather advisory: *Light to Moderate rainfall/thundershower is forecasted for today to Saturday in the block.* The day temperature will be 31-33°C and night temperature will be 24°C for upcoming 5 days. Maximum and minimum relative humidity will be 88-91% and 50-68% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Narasinghpur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	25/08/2026	26/08/2026	27/08/2026	28/08/2026	29/08/2026
Rainfall(mm)	8	7	6	14	25
T-Max(°C)	31	32	32	33	32
T-Min(°C)	24	24	24	24	24
Cloud Cover(octa)	7	6	7	8	8
RH Max (%)	91	89	88	89	91
RH Min (%)	60	63	58	50	68
Wind Speed (Kmph)	9	8	7	7	9
Wind Direction (deg)	198	255	244	189	145

Agromet Advisory

General advisory: Farmers should drain out excess water from vegetable and rice field. They may go for sowing, fertilizer, pesticide application, intercultural operations & human labour engagement before Friday based upon rainfall forecast.

Kharif Rice:

- The crop is at tillering stage. Maintain about 2-3 cm standing water during tillering stage. Problem of weeds may aggravate after transplanting, apply Bensulfuron methyl 0.6% + Pretilachlor 6% GR with sand mix at 1:1 ratio @ 4 kg/acre within 3-7 days after transplanting for controlling weeds.
- If leaf blast is observed in nursery, spray Tebuconazole 50% + Trifloxystrobin 25% WG @ 0.4 g or Isoprothiolane 40EC @ 1.5 ml per litre of water. Repeat the spray at 7-10 days interval.
- In Gall midge affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre
- Weekly Temperature below 28deg c and continuous rainfall lead to infestation of stem borer at tillering stage of rice, to control this apply Azadirachtin 0.15% EC @ 800 ml/acre or Cartap hydrochloride 4G @ 10kg/acre.
- Due to forecasted slight increase in temperature there may be increase in infestation of thrips, to control this Spray Lambda-cyhalothrin 5% EC @ 2 ml/ litre water or Thiamethoxam 25% WG @ 5g/ 15 litre water.
- Due to slight increase temperature and high humidity there will be chances of infestation of bacterial leaf streak, to control apply Plantomycin @ 1 g/litre along with Copper oxychloride @ 1g/litre of water using 200 litre of water per acre.

Maize: Maize crop is highly sensitive to waterlogged conditions during seedling stage. Hence, drain out excess water from the field as early as possible.

- After cessation of rains apply booster dose of fertilizer.
- Weeds may be controlled after rainfall.
- Stem fly in Maize infects the plants from sowing to 5 weeks after sowing, temperature between 25-30 deg. celsius and humidity above 60% favour pest incidence. In vegetative stage to control stem fly spray Methyl demeton @ 1 ml/ltr or Dimethoate @ 2ml/ltr or apply Carbofuran 3G granules @ 14 kg/acre.
- High humid conditions, temperature between 18-27 Deg. Celsius favours the occurrence of leaf blight in Maize. To control spray Azoxystrobin + Mancozeb 300 g /ac. Due to water logging condition stem rot may be found, then farmers should apply carbendazim @ 2g/ltr water.

VEGETABLES:

Anticipation of decrease in rainfall or less cloudy weather lead to infestation of mealy bug and mites in vegetables. Take up spraying of profenophos 50% EC @ 400ml/ac against mealy bug and propargite 57%EC/ fenazaquin 10%EC / Fenpyroximate 5% EC @ 400ml/ac against mites which are more prevalent in dry weather.

- Due to past rain, Chances of fruit rot infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary, spray second and third round at 15 days interval.
- If incidence of Damping off is noticed in Tomato, Brinjal, Chilli nurseries, drench nursery bed with Copper-Oxy-Chloride @ 3 g per liter of water.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever sucking pests incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. To prevent of soil borne diseases like seedling rot, stem rot and wilt diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control Cercospora leaf spot.

Fruits: Spray mancozeb 2g/litre, to control sigatoka leaf spot in banana. To control papaya yellow mosaic virus, spray Imidachloprid @ 0.5 ml / litre.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-646
Date- 25/08/2026

AGROMET ADVISORY BULLETIN FOR NIALI

District Past week average weather condition and Block past week average Rainfall condition

The block received average 10 mm and district received average 08 mm rainfall during the last week. The mean daily maximum and minimum temperature was 32.34°C (+0.34) and 23.62°C (-0.5), respectively. The mean relative humidity was 83-88%.

Forecasted period: Today(8.30IST) to 30th August (8.30IST)2026

Weather advisory: Light to Moderate rainfall/thundershower is forecasted for today to Saturday in the block. The day temperature will be 33°C and night temperature will be 25-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 37-90% and 49-68% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Niali (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	25/08/2026	26/08/2026	27/08/2026	28/08/2026	29/08/2026
Rainfall(mm)	8	7	6	14	24
T-Max(°C)	33	33	33	33	33
T-Min(°C)	25	26	26	26	26
Cloud Cover(octa)	7	5	7	7	7
RH Max (%)	37	37	80	80	90
RH Min (%)	60	60	56	49	68
Wind Speed(Kmph)	9	8	6	8	10
Wind Direction(deg)	168	243	215	176	138

Agromet Advisory

General advisory: Farmers should drain out excess water from vegetable and rice field. They may go for sowing, fertilizer, pesticide application, intercultural operations & human labour engagement before Friday based upon rainfall forecast.

Kharif Rice:

- The crop is at tillering stage. Maintain about 2-3 cm standing water during tillering stage. Problem of weeds may aggravate after transplanting, apply Bensulfuron methyl 0.6% + Pretilachlor 6% GR with sand mix at 1:1 ratio @ 4 kg/acre within 3-7 days after transplanting for controlling weeds.
- If leaf blast is observed in nursery, spray Tebuconazole 50% + Trifloxystrobin 25% WG @ 0.4 g or Isoprothiolane 40EC @ 1.5 ml per litre of water. Repeat the spray at 7-10 days interval.
- In Gall midge affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre
- Weekly Temperature below 28deg c and continuous rainfall lead to infestation of stem borer at tillering stage of rice, to control this apply Azadirachtin 0.15% EC @ 800 ml/acre or Cartap hydrochloride 4G @ 10kg/acre.
- Due to forecasted slight increase in temperature there may be increase in infestation of thrips, to control this Spray Lambda-cyhalothrin 5% EC @ 2 ml/ litre water or Thiamethoxam 25% WG @ 5g/ 15 litre water.
- Due to slight increase temperature and high humidity there will be chances of infestation of bacterial leaf streak, to control apply Plantomycin @ 1 g/litre along with Copper oxychloride @ 1g/litre of water using 200 litre of water per acre.

Sugarcane:

- Drain out excess water from the field as early as possible.
- After cessation of rains weeding and earthing up should be done in this time after that Top dressing of Urea @30 kg/acre for 30-60 days old sugarcane plant.
- For Red rot and wilt disease, spray the crop with Mancozeb 0.3% or carbendazim 0.1% or Azoxystrobin 18.2%SC/ Difenconazole 11.45SC @ 1ml per liter water in dry days.
- For early shoot borer Spray Chloropyriphos @ 400 ml/ac in 200 lts of water after cessation of rain.

VEGETABLES:

Anticipation of decrease in rainfall or less cloudy weather lead to infestation of Mealy bug and mites in vegetables. Take up spraying of profenophos 50% EC @ 400ml/ac against mealy bug and propargite 57%EC/ fenazaquin 10%EC / Fenpyroximate 5% EC @ 400ml/ac against mites which are more prevalent in dry weather.

- Due to past rain, Chances of fruit rot infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary, spray second and third round at 15 days interval.
- If incidence of Damping off is noticed in Tomato, Brinjal, Chilli nurseries, drench nursery bed with Copper-Oxy-Chloride @ 3 g per liter of water.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever sucking pests incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. To prevent of soil borne diseases like seedling rot, stem rot and wilt diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control Cercospora leaf spot.

Fruits: Spray mancozeb 2g/litre, to control sigatoka leaf spot in banana. To control papaya yellow mosaic virus, spray Imidachloprid @ 0.5 ml / litre.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-646
Date- 25/08/2026

AGROMET ADVISORY BULLETIN FOR NISCHINTAKOILI
District Past week average weather condition and Block past week average Rainfall condition

The block received average 07 mm and district received average 08 mm rainfall during the last week. The mean daily maximum and minimum temperature was 32.34°C (+0.34) and 23.62°C (-0.5), respectively. The mean relative humidity was 83-88%.

Forecasted period: Today(8.30IST) to 30th August (8.30IST)2026

Weatheradvisory: Light to Moderate rainfall/thundershower is forecasted for today to Saturday in the block. The day temperature will be 32-33°C and night temperature will be 25-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 37-90% and 49-68% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Nischintakoili (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	25/08/2026	26/08/2026	27/08/2026	28/08/2026	29/08/2026
Rainfall(mm)	8	7	6	14	24
T-Max(°C)	32	33	33	33	33
T-Min(°C)	25	26	26	26	26
Cloud Cover(octa)	7	6	6	7	8
RH Max (%)	37	37	80	80	90
RH Min (%)	60	60	56	49	68
Wind Speed (Kmph)	9	8	6	8	10
Wind Direction (deg)	168	243	215	176	138

Agromet Advisory

General advisory: Farmers should drain out excess water from vegetable and rice field. They may go for sowing, fertilizer, pesticide application, intercultural operations & human labour engagement before Friday based upon rainfall forecast.

Kharif Rice:

- The crop is at tillering stage. Maintain about 2-3 cm standing water during tillering stage. Problem of weeds may aggravate after transplanting, apply Bensulfuron methyl 0.6% + Pretilachlor 6% GR with sand mix at 1:1 ratio @ 4 kg/acre within 3-7 days after transplanting for controlling weeds.
- If **leaf blast** is observed in nursery, spray Tebuconazole 50% + Trifloxystrobin 25% WG @ 0.4 g or Isoprothiolane 40EC @ 1.5 ml per litre of water. Repeat the spray at 7-10 days interval.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre
- Weekly Temperature below 28deg c and continuous rainfall lead to infestation of **stem borer** at tillering stage of rice, to control this apply Azadirachtin 0.15% EC @ 800 ml/acre or Cartap hydrochloride 4G @ 10kg/acre.
- Due to forecasted slight increase in temperature there may be increase in infestation of **thrips**, to control this Spray Lambda-cyhalothrin 5% EC @ 2 ml/ litre water or Thiamethoxam 25% WG @ 5g/ 15 litre water.
- Due to slight increase temperature and high humidity there will be chances of infestation of **bacterial leaf streak**, to control apply Plantomycin @ 1 g/litre along with Copper oxychloride @ 1g/litre of water using 200 litre of water per acre.

Sugarcane:

- Drain out excess water from the field as early as possible.
- After cessation of rains weeding and earthing up should be done in this time after that Top dressing of Urea @30 kg/acre for 30-60 days old sugarcane plant.
- For **Red rot and wilt disease**, spray the crop with Mancozeb 0.3% or carbendazim 0.1% or Azoxystrobin 18.2%SC/ Difenconazole 11.45SC @ 1ml per liter water in dry days.
- For early shoot borer Spray Chloropyriphos @ 400 ml/ac in 200 lts of water after cessation of rain.

VEGETABLES:

Anticipation of decrease in rainfall or less cloudy weather lead to infestation of **Mealy bug** and **mites** in vegetables. Take up spraying of profenophos 50% EC @ 400ml/ac against mealy bug and propargite 57%EC/ fenazaquin 10%EC / Fenpyroximate 5% EC @ 400ml/ac against mites which are more prevalent in dry weather.

- Due to past rain, Chances of **fruit rot** infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary, spray second and third round at 15 days interval.
- If incidence of **Damping off** is noticed in Tomato, Brinjal, Chilli nurseries, drench nursery bed with Copper-Oxy-Chloride @ 3 g per liter of water.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever **sucking pests** incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. To prevent of soil borne diseases like **seedling rot, stem rot and wilt** diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control **Cercospora leaf spot**.

Fruits: Spray mancozeb 2g/litre, to control sigatoka leaf spot in banana. To control papaya yellow mosaic virus, spray Imidachloprid @ 0.5 ml / litre.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-646
Date- 25/08/2026

AGROMET ADVISORY BULLETIN FOR SALEPUR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 06 mm and district received average 08 mm rainfall during the last week. The mean daily maximum and minimum temperature was 32.34°C (+0.34) and 23.62°C (-0.5), respectively. The mean relative humidity was 83-88%.

Forecasted period: Today(8.30IST) to 30th August (8.30IST)2026

Weather advisory: *Light to Moderate rainfall/thundershower is forecasted for today to Saturday in the block.* The day temperature will be 33-34°C and night temperature will be 25°C for upcoming 5 days. Maximum and minimum relative humidity will be 37-90% and 49-68% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Salepur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	25/08/2026	26/08/2026	27/08/2026	28/08/2026	29/08/2026
Rainfall(mm)	8	7	6	15	24
T-Max(°C)	33	33	33	34	33
T-Min(°C)	25	25	25	25	25
Cloud Cover(octa)	7	6	7	8	8
RH Max (%)	37	37	80	80	90
RH Min (%)	60	60	56	49	68
Wind Speed (Kmph)	9	8	6	8	10
Wind Direction (deg)	168	243	215	176	138

Agromet Advisory

General advisory: Farmers should drain out excess water from vegetable and rice field. They may go for sowing, fertilizer, pesticide application, intercultural operations & human labour engagement before Friday based upon rainfall forecast.

Kharif Rice: The crop is at tillering stage. Maintain about 2-3 cm standing water during tillering stage. Problem of weeds may aggravate after transplanting, apply Bensulfuron methyl 0.6% + Pretilachlor 6% GR with sand mix at 1:1 ratio @ 4 kg/acre within 3-7 days after transplanting for controlling weeds.

• If **leaf blast** is observed in nursery, spray Tebuconazole 50% + Trifloxystrobin 25% WG @ 0.4 g or Isoprothiolane 40EC @ 1.5 ml per litre of water. Repeat the spray at 7-10 days interval.

• In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre

• Weekly Temperature below 28deg c and continuous rainfall lead to infestation of **stem borer** at tillering stage of rice, to control this apply Azadirachtin 0.15% EC @ 800 ml/acre or Cartap hydrochloride 4G @ 10kg/acre.

• Due to forecasted slight increase in temperature there may be increase in infestation of **thrips**, to control this Spray Lambda-cyhalothrin 5% EC @ 2 ml/ litre water or Thiamethoxam 25% WG @ 5g/ 15 litre water.

• Due to slight increase temperature and high humidity there will be chances of infestation of **bacterial leaf streak**, to control apply Plantomycin @ 1 g/litre along with Copper oxychloride @ 1g/litre of water using 200 litre of water per acre.

Maize: Maize crop is highly sensitive to waterlogged conditions during seedling stage. Hence, drain out excess water from the field as early as possible.

• After cessation of rains apply booster dose of fertilizer.

• Weeds may be controlled after rainfall.

• **Stem fly** in Maize infects the plants from sowing to 5 weeks after sowing, temperature between 25-30 deg. celsius and humidity above 60% favour pest incidence. In vegetative stage to control stem fly spray Methyl demeton @ 1 ml/lit or Dimethoate @ 2ml/lit or apply Carbofuran 3G granules @ 14 kg/acre.

• High humid conditions, temperature between 18-27 Deg. Celsius favours the occurrence of **leaf blight** in Maize. To control spray Azoxystrobin + Mancozeb 300 g /ac. Due to water logging condition stem rot may be found, then farmers should apply carbendazim @ 2g/ltr water.

Jute: Recent rain has aggravated **weed problem** in early sown crop. To control grass weeds, apply Quisalofof-pethyl @ 2ml/litre of water at 2-3 leaf stage of weeds. After rain in vegetative stage to control **stem rot of jute**, spray copper oxychloride @ 3-4 gram/liter of water.

VEGETABLES:

Anticipation of decrease in rainfall or less cloudy weather lead to infestation of **Mealy bug** and **mites** in vegetables. Take up spraying of profenophos 50% EC @ 400ml/ac against mealy bug and propargite 57% EC / fenazaquin 10% EC / Fenpyroximate 5% EC @ 400ml/ac against mites which are more prevalent in dry weather.

• Due to past rain, Chances of **fruit rot** infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary, spray second and third round at 15 days interval.

• If incidence of **Damping off** is noticed in Tomato, Brinjal, Chilli nurseries, drench nursery bed with Copper-Oxy-Chloride @ 3 g per liter of water.

• Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever **sucking pests** incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. To prevent of soil borne diseases like **seedling rot, stem rot and wilt** diseases apply 3g of Copper oxychloride per litre of water around the main stem.

• Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control **Cercospora leaf spot**.

Fruits: Spray mancozeb 2g/litre, to control sigatoka leaf spot in banana. To control papaya yellow mosaic virus, spray Imidachloprid @ 0.5 ml / litre.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-646
Date- 25/08/2026

AGROMET ADVISORY BULLETIN FOR TANGI-CHOUDWAR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 12 mm and district received average 08 mm rainfall during the last week. The mean daily maximum and minimum temperature was 32.34°C (+0.34) and 23.62°C (-0.5), respectively. The mean relative humidity was 83-88%.

Forecasted period: Today(8.30IST) to 30th August (8.30IST)2026

Weather advisory: *Light to Moderate rainfall/thundershower is forecasted for today to Saturday in the block.* The day temperature will be 33°C and night temperature will be 25°C for upcoming 5 days. Maximum and minimum relative humidity will be 37-90% and 49-68% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Tangi-Choudwar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	25/08/2026	26/08/2026	27/08/2026	28/08/2026	29/08/2026
Rainfall(mm)	8	7	6	15	24
T-Max(°C)	33	33	33	33	33
T-Min(°C)	25	25	25	25	25
Cloud Cover(octa)	7	6	6	8	8
RH Max (%)	37	37	80	80	90
RH Min (%)	60	60	56	49	68
Wind Speed (Kmph)	9	8	6	8	10
Wind Direction (deg)	168	243	215	176	138

Agromet Advisory

General advisory: Farmers should drain out excess water from vegetable and rice field. They may go for sowing, fertilizer, pesticide application, intercultural operations & human labour engagement before Friday based upon rainfall forecast.

Kharif Rice:

- The crop is at tillering stage. Maintain about 2-3 cm standing water during tillering stage. Problem of weeds may aggravate after transplanting, apply Bensulfuron methyl 0.6% + Pretilachlor 6% GR with sand mix at 1:1 ratio @ 4 kg/acre within 3-7 days after transplanting for controlling weeds.
- If **leaf blast** is observed in nursery, spray Tebuconazole 50% + Trifloxystrobin 25% WG @ 0.4 g or Isoprothiolane 40EC @ 1.5 ml per litre of water. Repeat the spray at 7-10 days interval.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre
- Weekly Temperature below 28deg c and continuous rainfall lead to infestation of **stem borer** at tillering stage of rice, to control this apply Azadirachtin 0.15% EC @ 800 ml/acre or Cartap hydrochloride 4G @ 10kg/acre.
- Due to forecasted slight increase in temperature there may be increase in infestation of **thrips**, to control this Spray Lambda-cyhalothrin 5% EC @ 2 ml/ litre water or Thiamethoxam 25% WG @ 5g/ 15 litre water.
- Due to slight increase temperature and high humidity there will be chances of infestation of **bacterial leaf streak**, to control apply Plantomycin @ 1 g/litre along with Copper oxychloride @ 1g/litre of water using 200 litre of water per acre.

Maize: Maize crop is highly sensitive to waterlogged conditions during seedling stage. Hence, drain out excess water from the field as early as possible.

- After cessation of rains apply booster dose of fertilizer.
- Weeds may be controlled after rainfall.
- Stem fly** in Maize infects the plants from sowing to 5 weeks after sowing, temperature between 25-30 deg. celsius and humidity above 60% favour pest incidence. In vegetative stage to control stem fly spray Methyl demeton @ 1 ml/ltr or Dimethoate @ 2ml/ltr or apply Carbofuran 3G granuals @ 14 kg/acre.
- High humid conditions, temperature between 18-27 Deg. Celsius favours the occurrence of **leaf blight** in Maize. To control spray Azoxystrobin + Mancozeb 300 g /ac. Due to water logging condition stem rot may be found, then farmers should apply carbendazim @ 2g/ltr water.

VEGETABLES:

Anticipation of decrease in rainfall or less cloudy weather lead to infestation of **Mealy bug** and **mites** in vegetables. Take up spraying of profenophos 50% EC @ 400ml/ac against mealy bug and propargite 57%EC/ fenazaquin 10%EC / Fenpyroximate 5% EC @ 400ml/ac against mites which are more prevalent in dry weather.

- Due to past rain, Chances of **fruit rot** infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary, spray second and third round at 15 days interval.
- If incidence of **Damping off** is noticed in Tomato, Brinjal, Chilli nurseries, drench nursery bed with Copper-Oxy-Chloride @ 3 g per liter of water.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever **sucking pests** incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. To prevent of soil borne diseases like **seedling rot, stem rot and wilt** diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control **Cercospora leaf spot**.

Fruits: Spray mancozeb 2g/litre, to control sigatoka leaf spot in banana. To control papaya yellow mosaic virus, spray Imidachloprid @ 0.5 ml / litre.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-646
Date- 25/08/2026

AGROMET ADVISORY BULLETIN FOR TIGIRIA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 12 mm and district received average 08 mm rainfall during the last week. The mean daily maximum and minimum temperature was 32.34°C (+0.34) and 23.62°C (-0.5), respectively. The mean relative humidity was 83-88%.

Forecasted period: Today(8.30IST) to 30th August (8.30IST)2026

Weather advisory: Light to Moderate rainfall/thundershower is forecasted for today to Saturday in the block. The day temperature will be 32-33°C and night temperature will be 25°C for upcoming 5 days. Maximum and minimum relative humidity will be 37-90% and 49-68% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Tigiria (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	25/08/2026	26/08/2026	27/08/2026	28/08/2026	29/08/2026
Rainfall(mm)	7	7	6	15	26
T-Max(°C)	32	32	33	33	32
T-Min(°C)	25	25	25	25	25
Cloud Cover(octa)	7	5	6	8	8
RH Max (%)	37	37	80	80	90
RH Min (%)	60	60	56	49	68
Wind Speed (Kmph)	9	8	6	8	10
Wind Direction (deg)	168	243	215	176	138

Agromet Advisory

General advisory: Farmers should drain out excess water from vegetable and rice field. They may go for sowing, fertilizer, pesticide application, intercultural operations & human labour engagement before Friday based upon rainfall forecast.

Kharif Rice:

- The crop is at tillering stage. Maintain about 2-3 cm standing water during tillering stage. Problem of weeds may aggravate after transplanting, apply Bensulfuron methyl 0.6% + Pretilachlor 6% GR with sand mix at 1:1 ratio @ 4 kg/acre within 3-7 days after transplanting for controlling weeds.
- If **leaf blast** is observed in nursery, spray Tebuconazole 50% + Trifloxystrobin 25% WG @ 0.4 g or Isoprothiolane 40EC @ 1.5 ml per litre of water. Repeat the spray at 7-10 days interval.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre
- Weekly Temperature below 28deg c and continuous rainfall lead to infestation of **stem borer** at tillering stage of rice, to control this apply Azadirachtin 0.15% EC @ 800 ml/acre or Cartap hydrochloride 4G @ 10kg/acre.
- Due to forecasted slight increase in temperature there may be increase in infestation of **thrips**, to control this Spray Lambda-cyhalothrin 5% EC @ 2 ml/ litre water or Thiamethoxam 25% WG @ 5g/ 15 litre water.
- Due to slight increase temperature and high humidity there will be chances of infestation of **bacterial leaf streak**, to control apply Plantomycin @ 1 g/litre along with Copper oxychloride @ 1g/litre of water using 200 litre of water per acre.

Maize: Maize crop is highly sensitive to waterlogged conditions during seedling stage. Hence, drain out excess water from the field as early as possible.

- After cessation of rains apply booster dose of fertilizer.
- Weeds may be controlled after rainfall.
- Stem fly** in Maize infects the plants from sowing to 5 weeks after sowing, temperature between 25-30 deg. celsius and humidity above 60% favour pest incidence. In vegetative stage to control stem fly spray Methyl demeton @ 1 ml/lit or Dimethoate @ 2ml/lit or apply Carbofuran 3G granules @ 14 kg/acre.
- High humid conditions, temperature between 18-27 Deg. Celsius favours the occurrence of **leaf blight** in Maize. To control spray Azoxystrobin + Mancozeb 300 g /ac. Due to water logging condition stem rot may be found, then farmers should apply carbendazim @ 2g/ltr water.

VEGETABLES:

Anticipation of decrease in rainfall or less cloudy weather lead to infestation of **Mealy bug and **mites** in vegetables. Take up spraying of profenophos 50% EC @ 400ml/ac against mealy bug and propargite 57%EC/ fenazaquin 10%EC / Fenpyroximate 5% EC @ 400ml/ac against mites which are more prevalent in dry weather.**

- Due to past rain, Chances of **fruit rot** infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary, spray second and third round at 15 days interval.
- If incidence of **Damping off** is noticed in Tomato, Brinjal, Chilli nurseries, drench nursery bed with Copper-Oxy-Chloride @ 3 g per liter of water.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever **sucking pests** incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. To prevent of soil borne diseases like **seedling rot, stem rot and wilt** diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control **Cercospora leaf spot**.

Fruits: Spray mancozeb 2g/litre, to control sigatoka leaf spot in **banana**. To control **papaya yellow mosaic virus**, spray Imidachloprid @ 0.5 ml / litre.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area






GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-646
Date- 25/08/2026

				
MEGHDOOT APP	DAMINI APP	FACEBOOK (Damu Cuttack)	BLOCK SPECIFIC WHAT'S APP GROUPS	@DAMUCUTTACK

- Kindly install and use **“Meghdoot mobile App”** for Cuttack district weather forecast, warnings and weather based agromet advisory services.
<https://play.google.com/store/apps/details?id=com.aas.meghdoot>
- Kindly install and use **“Damini mobile App”** for location specific for lightening forecast and warnings.
<https://play.google.com/store/apps/details?id=com.lightening.live.damini>
- Follow the **block specific whats app groups** and join in your block wise groups for getting weather based agromet advisory services twice in a week (Tuesday and Friday) and other weather warning based farm management advisories.
- Follow the **Official Website links** to get the weather based agromet advisory service bulletins (English & Odia) twice in a week (Tuesday and Friday).

1	ICAR-NRRI, Cuttack, Odisha	https://icar-nrri.in/agromate/	
2	Agrimet division, IMD, Pune	https://imdagrimet.gov.in/imdproject/AGIndex.php	



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

